

Responses to Cortical Spreading Depression under Oxygen Deficiency

J. Sonn* and A. Mayevsky

The Mina & Everard Goodman, Faculty of Life Sciences and Leslie and Susan Gonda Multidisciplinary Brain Research Center, Bar-Ilan University RAMAT-GAN 52900, Israel

Abstract: *Objectives:* The effect of cortical spreading depression (CSD) on extracellular K^+ concentrations ($[K^+]_e$), cerebral blood flow (CBF), mitochondrial NADH redox state and direct current (DC) potential was studied during normoxia and three pathological conditions: hypoxia, after NOS inhibition by L-NAME and partial ischemia.

Methods: A special device (MPA) was used for monitoring CSD wave propagation, containing: mitochondrial NADH redox state and reflected light, by a fluorometry technique; DC potential by Ag/AgCl electrodes; CBF by laser Doppler flowmetry; and $[K^+]_e$ by a mini-electrode.

Results and Discussion: 1. CSD under the 3 pathological conditions caused an initial increase in NADH and a further decrease in CBF during the first phase of CSD, indicating an imbalance between oxygen supply and demand as a result of the increase in oxygen requirements. 2. The hyperperfusion phase in CBF was significantly reduced during hypoxia and ischemia showing a further decline in oxygen supply during CSD. 3. CSD wave duration increased during the pathological conditions, showing a disturbance in energy production. 4. Extracellular K^+ levels during CSD, increased to identical levels during normoxia and during the three pathological groups, indicating correspondingly increase in oxygen demand. 5. The special design of the MPA enabled identifying differences in the simultaneous responses of the measured parameters, which may indicate changes in the interrelation between oxygen demand, oxygen supply and oxygen balance during CSD propagation, under the conditions tested. 6. In conclusion, brain oxygenation was found to be a critical factor in the responses of the brain to CSD.

Keywords: Brain oxygenation, cerebral blood flow, extracellular K^+ , hypoxia, mitochondrial NADH, nitric oxide synthase inhibition, partial ischemia.

INTRODUCTION

Cortical spreading depression (CSD) is a transient neuronal depolarization that is accompanied by a negative shift in the DC potential and a flattening in EEG [1], changes in membrane permeability as well as in the ionic homeostasis [2], thereby increasing energy consumption [3] and blood flow [4, 5]. The activation of ion pumps, following the changes in extracellular and intracellular ion levels consumes energy [5, 6]. Therefore, the responses of the brain to CSD were used as a tool that increases oxygen demand [7] and serves as an indicator to the tissue energetic-metabolic state [7-9]. In the case of healthy normoxic brain, repeated CSD waves do not cause any damage to the tissue [10]. In contradiction, CSD is thought to be involved in the mechanism of the neurological symptoms in migraine with aura [4, 11] and it has also been reported that CSD may occur spontaneously under hypoxic or ischemic conditions [5, 11-13], following traumatic brain injury [2, 14, 15] and was also found to appear after epileptic seizures [11, 16].

The brain is completely dependent upon a continuous supply of blood flow providing oxygen, glucose, nutrients etc. Therefore, during pathological conditions, where oxygen supply is limited, such as hypoxia or ischemia, any increase in oxygen demand will cause an oxygen deficiency that will end with severe functional disorders and irreversible brain

damage [17]. Under such conditions, CSD will increase tissue oxygen demand and lead to a further increase in oxygen deficiency [7, 18] which will augment tissue damage [12, 19, 20].

Nitric oxide (NO) acts as a neurotransmitter [21] as well as a regulator of cerebral blood flow [22, 23]. NO generation is closely linked with excessive glutamate release and has been implicated in ischemic and traumatic neurodegeneration [22, 24]. However, a decline in CBF or in oxygen supply (i.e. under ischemia and hypoxia) was found to disturb NO production [25, 26]. Studies showed that the increase in CBF, during CSD after NO synthase inhibition, was diminished [27, 28]. In our previous studies we found that under normal oxygen supply, increasing oxygen demand by CSD induction, caused an increase in CBF in parallel to oxidation cycles in mitochondrial NADH [7, 8, 29]. We have also shown that responses (qualitative) to CSD during hypoxia and ischemia were controlled by the limited level of blood supply and low blood oxygen concentration. Under these conditions no oxidation waves were observed in NADH [7].

The aims of the present study were: 1. To find the interrelation between the changes in extracellular K^+ (representing oxygen demand), CBF (representing oxygen supply) and mitochondrial NADH (representing oxygen balance) during CSD propagation, under hypoxia, ischemia and after NOS inhibition. 2. To examine the quantitative responses to CSD under hypoxia, ischemia, after NOS inhibition, compared to CSD under normoxic conditions.

*Address correspondence to this author at the Faculty of Life Sciences Bar-Ilan University RAMAT-GAN 52900, Israel; Tel: 972-3-5317558; Fax: 972-3-5352184; E-mail: sonnju1@gmail.com

compare the order (in time) of amplitude changes (maximum, minimum) in the measured parameters according to the beginning of the CSD wave propagation. Thus, we were able to evaluate the interrelation between the measured parameters during the CSD wave propagation.

GENERAL EXPERIMENTAL PROCEDURE

Animal Preparation

The Institutional Animal Care and Use Committee of Bar Ilan University approved all experimental protocols. Male Wistar rats weighing 200-250 g were anesthetized with a 0.3 ml/100g IP injection of Equithesin (each ml containing 42.51 mg chloral hydrate, 9.72 mg pentobarbital, 21.25 mg magnesium sulfate, 44.43% w/v propylene glycol, 11.5% alcohol and water). A thermistor probe (Yellow Springs Instruments Co. Inc., type 402, Ohio, USA) was inserted into the rectum to measure and control body temperature during surgery and experiment. The cranium was exposed and a thin rectangular Plexiglas pattern was located on it. The pattern contained two scratches: the MPA probe (6.4×3.3 mm) and the KCl push-pull cannula (2 mm diameter) in the exact shape as it is schematically illustrated in Fig. (1). Two holes were drilled: one in the frontal bone for the KCl cannula and the other in the ipsilateral parietal bone for the MPA implantation. No harm was done to the sagittal sinus during MPA implantation. The dura mater was gently removed only in the monitoring site and not where CSD was initiated. The push-pull cannula enabled flushing the brain with diluted KCl solution (0.5M) in order to induce CSD waves. If 0.5M did not induce CSD waves, then the concentration of KCl solution was increased to 1M up to saturated KCl solution (in increments of 0.5M) until the initiation of CSD waves. Two other holes were drilled for screws located in the contra lateral hemisphere that helped to securely hold the system to the cranium. All components were fixated to the skull using dental acrylic cement. About 15 minutes later, the animals were inhaled with pure N₂ for about 20 seconds in order to check maximum NADH increase under anoxic conditions. This data was used to ensure no brain damage during the surgery procedure. The experiment started about 60-90 minutes after the surgery (after ensuring complete recovery from the surgery procedure). The animals were kept lightly anesthetized (leg reflex still remained and body temperature did not decrease under 36.5°C) by continuous IP infusion of 0.085-0.17 ml/h Equithesin (Eth.) depending on the rat's weight. This level of anesthesia also prevented spontaneous CSD waves. At the end of the experiment the animals were deeply reanesthetized and sacrificed by N₂ inhalation. In the hypoxia group the femoral artery was cannulated and arterial blood samples were withdrawn for pH, pO₂ and pCO₂ analysis.

The decrease of oxygen delivery to the brain tissue was achieved by three different perturbations using three different animal protocols: hypoxia, partial ischemia and by inhibition of nitric oxide (NO) synthesis with N^ω-nitro-L-arginine methyl ester (C₇H₁₅N₅O₄.HCl, L-NAME).

Hypoxia was induced by inhalation of a gas mixture containing 12% O₂ + 88% N₂ for 17-20 min. Partial ischemia was induced by permanent bilateral carotid artery occlusion 24 hours before monitoring.

Hypoxia Protocol

After recovery from the surgery procedure three CSD waves were induced with an interval of one hour between each wave. The first wave was induced under normoxic conditions (control); the second, under hypoxic conditions and the third under normoxic conditions. The CSD wave under hypoxic conditions was induced during gas mixture inhalation, when the parameters showed a new steady state (after about 6 minutes of inhalation). The hypoxic period continued for about 17-20 minutes until a recovery in CSD wave was achieved. The third CSD wave was induced one hour after the hypoxic CSD wave, under normoxic conditions, to verify recovery from hypoxia. This group contained 13 rats. This protocol contained 2 control groups: one group that passed the same protocol under normoxic conditions, without inducing hypoxia, (n=11) and another control group of rats that were inhaled with the gas mixture (in the same range of time) and no CSD waves were induced. This control group contained 8 rats and was done in order to ensure the effect of hypoxia on the measured parameters devoid of the effect of CSD propagation. Arterial blood samples were taken during normoxia and hypoxia.

Protocol of inhibition of Nitric Oxide Synthase (NOS)

Inhibition of nitric oxide synthesis (n=10) was achieved by IP injection of 50 mg/kg L-NAME solution. The protocol contained 3 CSD waves. The first CSD wave was induced under normoxia-control conditions. One hour later, 50 mg/kg of L-NAME was injected IP. Two additional CSD waves were induced one and two hours after L-NAME injection. The experiment and the monitoring period continued 1.5 hours after the last initiation of CSD wave. In another group of rats (n=11) 4 CSD waves were induced with an interval of one hour between each wave under normoxic conditions. This group of rats was considered as a control group for hypoxia and L-NAME groups.

Ischemia Protocol

Partial ischemia (n=9) was induced by permanent bilateral carotid artery occlusion. Twenty-four hours later the rats were reanesthetized and underwent preparation for the experiment and monitoring. Sham rats (n=8) underwent the same experimental procedure without carotids occlusion. CSD waves were induced (by epidural application of 0.5-2M KCl solution, using the lowest concentration that induced CSD), in lightly anesthetized rats in the partial ischemic and in sham rats.

Data Acquisition

The NADH fluorescence, reflected light, CBF, [K⁺]_e and DC steady potential that were monitored by the MPA system were recorded continuously, simultaneously and in real time using Coda software for a 16-channel computerized acquisition and storage system (DATAQ Instruments, MN, USA).

Data Analysis and Calculation

Various time and amplitude values were calculated during the CSD wave cycles. In the present article we will present: The maximum (VT_{max}) and minimum (VT_{min}) amplitude values, as well as wave duration that were calculated for every parameter recorded by the MPA system. Wave duration was calculated as the time it took the CSD wave to

Fig. (5). The effect CSD wave propagation on maximum (VTmax-black bars) and minimum (VTmin-gray bars) amplitudes of reflected light (reflectance), cerebral blood flow (CBF) and mitochondrial NADH redox state under normoxic (control) and after a single dose of L-NAME injected IP. L-NAME1 – CSD induced one hour after L-NAME injection and L-NAME2, CSD induced 2 hours after L-NAME injection. Arrows sign the initial reduction wave in NADH (grid bar) recorded during CSD. Data are presented as mean \pm S.E. * $p < 0.05$, ** $p < 0.005$, *** $p < 0.0005$ CSD during normoxia vs. CSD after L-NAME.

pared all changes to it. In addition, spontaneous oscillations were seen during ischemia, in CBF, reflectance and in NADH during the recordings, in an identical rate as seen during hypoxia and after L-NAME administration. The oscillations in mitochondrial NADH were found parallel and in an opposite direction to CBF.

Inductions of CSDs under ischemic conditions caused 3 kinds of variable responses (in different rats) in the measured parameters and are demonstrated in Fig. (6). Fig. (6A) shows that during CSD NADH increased, CBF decreased first, and only later, increased slightly (during the recovery phase). Extracellular K^+ increased and DC potential decreased. Trace 6B shows a large increase in reflectance followed by a small decrease, an increase followed by a decrease in NADH redox state, and a clear initial deep decrease in CBF, that was followed by an increase. Trace 6C shows an increase in NADH and in CBF. In all the ischemic traces (Fig. 6) reflectance changed in a biphasic manner, an increase that was followed by a decrease and the time to recovery from CSD was prolonged (in all the measured parameters) as compared to normoxic conditions (for example Figs. 2A, 4A).

Fig. (7) shows the mean \pm S.E. % values of reflectance, CBF and NADH during CSD induced in sham and ischemic brains. The increases in $[K^+]_e$ levels were up to 18.38 ± 1.51 mM and 19.31 ± 2.74 mM and the decreases in DC

potential were of 13.6 ± 0.5 mV and 12.86 ± 0.56 mV in sham and ischemic rats, respectively. No significant differences were found in $[K^+]_e$ and in DC potential between sham and ischemic rats. In the ischemic experiments, reflectance changed in a biphasic pattern as during normoxia but the increase phase was significantly higher ($p < 0.005$) during ischemia indicating a greater decrease in tissue blood volume during CSD under ischemic conditions as compared to sham.

CBF during CSD under normoxic conditions (sham) showed a decline that was followed by a large increase whereas, during ischemia the initial decrease was significantly larger ($p < 0.005$) and the increase phase was significantly smaller ($p < 0.05$).

NADH during CSD changed in a biphasic pattern in the sham rats: an oxidation wave was followed by a reduction one. Whereas, in the ischemic brain an initial and significant ($p < 0.0005$) reduction of about 30% was found before the biphasic pattern. Furthermore, the oxidation wave was significantly smaller ($p < 0.05$) and so was the late reduction cycle ($p < 0.0005$).

CSD wave durations under hypoxia, ischemia and one hour after L-NAME administration are summarized in Table 1. The data shows that the recovery from CSD prolonged in most measured parameters, during these pathological conditions, where oxygen availability to tissue was limited.




Fig. (6). Analog tracings presenting the effect of CSD on the measured parameters of three (A, B and C) different partial ischemic brains. Abbreviations as explained in Figs. (2 and 4).




Fig. (7). The effect of CSD wave propagation on maximum (VTmax-black bars) and minimum (VTmin-gray bars) amplitudes of reflected light (reflectance), cerebral blood flow (CBF) and mitochondrial NADH redox state on normoxic (sham) and partial ischemic brains. Arrows sign the initial reduction wave in NADH (grid bar) recorded during CSD. Data are presented as mean \pm S.E. * p<0.05, ** p<0.005, *** p<0.0005 of CSD on sham brains vs. CSD on partial ischemic brains.

The Interrelation Between CBF and NADH Redox State During CSD

Extracellular $[K^+]$ during CSD increased similarly during all conditions tested (no statistical differences were found).

In contradiction, the simultaneous measurements of CBF and NADH during CSD propagation, showed different trends, compared to normoxic conditions, and illustrated in Figs. (2, 4 and 6) by dotted lines. The parallel changes in CBF and

NADH during CSD under normoxic conditions (Figs. 2A and 4A) were totally different as compared to the three other conditions (Figs. 2B, 4B, 6A, 6B, 6C): under normoxia the oxidation in NADH preceded the maximum increase in CBF (Figs. 2A, 4A), whereas during hypoxia, ischemia and after L-NAME administration an initial reduction cycle in NADH was seen in parallel to the initial decrease in CBF (first dotted line in 2B, 4B, 6A, 6B, 6C) which raised during these pathological conditions. The increase in CBF during CSD (second dotted line in Figs. 2B, 4B, 6A, 6B, 6C) was observed afterwards. Parallel to the increase in CBF a decrease in NADH (oxidation) was observed. These results can demonstrate a different interrelation between CBF and NADH during CSD under these pathological conditions compared to control.

DISCUSSION

The results in the current study show that induction of CSD caused an increase in extracellular K^+ that reached identical levels under all perturbations tested, showing that the increase in oxygen demand by inducing CSD reached equal levels under normoxia, hypoxia, ischemia and after L-NAME administration (Figs. 2, 4, 6 and data (Mean \pm S.E) reported in pages 5, 6, 7). In healthy and normal tissue, these changes in ion homeostasis are associated with CBF and oxygen delivery elevation, leaving the tissue intact for many hours [10]. But for tissue in which blood flow or oxygen delivery is disturbed, compensation will be reliable on the availability of oxygen. Furthermore, increasing oxygen demand under such circumstances can cause a chain of events that will enhance or lead to brain damage: such as disruption in ion homeostasis, acidosis, release of glutamate and excitatory amino acids (EAs) etc [11, 19].

Brain Responses to NO Synthase Inhibition

We found that L-NAME injection decreased CBF levels, as was found by previous studies investigating the effect of NOS inhibition [22, 33-35]. In parallel, we found an increase in mitochondrial NADH showing a decline in oxygen delivery to the brain tissue which may reduce tissue oxygen balance. In addition, some studies [22, 33, 36] showed that L-NAME did not cause changes in arterial blood gases. Macrae showed [36] that NOS inhibition by L-NAME caused a decline in CBF in different brain areas which was not associated by a reduction in glucose consumption. Also Iadecola *et al.* in their review article [22] showed that oxygen and glucose consumption did not change as a result of NOS inhibition. Furthermore, Brown [37, 38] hypothesized and showed that NO is responsible to mitochondrial respiration regulation by inhibiting cytochrome oxidase. Therefore, the inhibition of NOS can lead to an increase in oxygen consumption [38]. Kurzelewski, *et al.*, [39] found, in isolated rat heart, that inhibition of NOS resulted in an increase in oxygen consumption but did not affect glucose or FFA oxidation. Our results showed an increase in mitochondrial NADH redox state which can result from the increase in oxygen consumption due to NOS inhibition by L-NAME. Besides, the elevation in extracellular K^+ can indicate that Na^+-K^+ -ATPase turnover decreased, which can signify a decrease in energy production. In addition, NADH elevation may also occur due to the decline in CBF which reduced oxygen delivery and lead to ATP depletion in the brain tissue.

Responses to CSD Under NO Synthase Inhibition

NO is known to be involved in regulation of CBF following neuronal activation [40]. In addition, it was shown that the mechanism of CSD propagation involves activation of glutamate and NMDA receptors [41] which causes post-synaptic calcium influx triggering NOS activation [22, 34, 42]. Thus, the question is whether NO is involved in the noticed vascular changes and in CBF regulation during CSD propagation? The early increase in reflectance and the decrease in CBF (CSD during the second hour after L-NAME administration) before the large increase (Figs. 4, 5) could result from the vasoconstriction effect due to the increase in extracellular K^+ concentrations (Fig. 4) resulted from neuronal depolarization following CSD propagation. A brief similar decline in CBF was seen by others [33, 34, 36] showing that NO could play an important vasodilative role during the early phase of CSD. The brief decline in CBF consequences an initial increase in reflectance and reduction in NADH (Figs. 4, 5) showing a decrease in blood volume and oxygen delivery during the brief phase of CSD after L-NAME administration. During the second phase of CSD, oxidation cycles were observed in NADH, parallel to a large increase in CBF that did not differ in amplitude changes from the control CSDs. These results indicate that other mediators than NO take a role in the vasodilatation mechanism during the hyperperfusion phase of CSD. Similar [33, 43, 44] and different [45, 46] results were found by others, where Akgoren and Dirnagl groups concluded that NO plays an important role in the coupling mechanism between CBF and neuronal activation. In addition, in rabbits and cats the CSD hyperperfusion phase decreased after NOS inhibition [27, 28, 47] which can indicate that the effect of NO on CBF changes during CSD may depend on the species investigated. However, although we found an increase in CBF during the second phase of CSD, the increase in wave duration (Table 1) may indicate that inhibition of NOS leads to energy depletion, which prolongs the brain tissue ability to recover from an increase in oxygen demand caused by CSD.

Brain Responses to Hypoxia

Systemic hypoxia decreased arterial blood pressure and arterial pO_2 . The reduction in mean arterial blood pressure was found similar to the "hypoxia normotension" conditions reported by Sukhotinsky *et al.*, [48]. In addition, the decline in arterial pO_2 stimulated the production of a wide variety of vasodilator metabolites, such as adenosine, NO, Ca^{2+} , H^+ and K^+ levels, prostaglandins and others, causing an increase in CBF [49]. It seems that the concomitant increase in CBF did not succeed completely to compensate the brain's tissue oxygen demand and as a result, a significant increase in mitochondrial NADH was noticed (see results section page 4). In the review article of Mayevsky and Rogatsky [9], the authors showed that mitochondrial NADH increased in an inversely correlation to the decrease in FiO_2 levels.

Responses to CSD Under Hypoxia

The initial phase, during CSD under systemic hypoxia, showed a brief decline in oxygen supply (CBF decreased and reflectance increased) and in oxygen balance (initial reduction in NADH) (Figs. 2, 3). Identical results were shown by others [48]. This initial decline in CBF can result from the increase in extracellular K^+ levels that is known to affect

vasoconstriction [2, 7, 50, 51], and consequently an added initial reduced phase (increase) in NADH redox state (Fig. 2B). In addition, tissue hypoxia can increase NO levels which may inhibit mitochondrial respiration [37, 38, 52] causing an additional increase in NADH and deplete ATP levels. Additionally, the increased phase in CBF (hyperperfusion phase) was also diminished significantly (Figs. 2B, 3) which can result from the initial high levels of vasoactive substances already present in the microcirculatory bed (resulted from the hypoxia), and therefore the hemodynamic responses to CSD are reduced. This decline in oxygen supply and in ATP levels prolonged wave duration and slowed down the recovery time from CSD (Table 1) during hypoxia. Similar results were shown in our previous studies [7, 53] as well identical changes in CBF, DC potential and in wave duration were reported by others [48].

Responses to CSD Under Partial Ischemia

We used the bilateral common carotid arteries occlusion as a model for partial ischemia. Several works on rats and gerbils showed that occlusion of one or both carotid arteries caused a decrease in CBF and an increase in NADH [54, 55]. The increase in NADH was found proportional to the decline in CBF [55] and was accompanied by changes in ionic homeostasis [2, 55-57].

Our data shows that induction of CSD waves on partial ischemic brains resulted in about 3 types of variable changes in the measured parameters, which were expressed in the amplitude levels of reflectance, CBF and NADH (Fig. 6). The variable reactions to CSD under ischemia could result from a variability in the intensity of brain tissue damage resulted from the bilateral carotid occlusion.

The initial increase in reflectance, the initial decline in CBF and the early increase (reduction cycle) in NADH fluorescence (Figs. 6, 7) can indicate a reduction in blood flow and volume (reflectance) and a decline in oxygen delivery to the ischemic brain tissue, during the early phase of CSD. The decline in CBF can also be explained as a result of microcirculatory vasoconstriction caused by the increase in extracellular K^+ levels. Identical results during CSD were shown under hypoxic conditions (Figs. 2, 3), in our previous study [7] and by others [48]. In addition, we showed [53] a decrease in tissue HbO_2 parallel to the increase in NADH. Also Back *et al.*, [58] showed a further decrease in tissue pO_2 and no changes in CBF during spontaneous CSDs after Middle Cerebral Artery occlusion. Besides, our data shows that the hyperperfusion phase during CSD decreased, and the oxidation wave in NADH almost disappeared. These findings indicate that increasing oxygen demand (by CSD) in the ischemic brain, limits tissue abilities to compensate these deficiencies leading to a reduction in Na^+K^+ ATPase pump turnover, which prolongs the recovery from CSDs (Tables 1). Similar responses in wave duration were reported by others [58] and by our group [7, 59].

The Interrelation Between CBF and NADH Redox State During CSD

Our data indicate that inhibition of NO synthesis, hypoxia or partial ischemia resulted in a decline in energy production, caused by a disruption in the balance between O_2 supply and O_2 demand (Figs. 2-7). The metabolic and hemo-

dynamic oscillations that were observed after induction of each of the tested pathological conditions, suggest a dynamic linkage between metabolic and vascular processes, which can indicate a tight coupling between oxygen supply (CBF) and energy balance (NADH). Identical observations, during partial ischemia, after L-NAME injection, after CO poisoning and brain trauma, were found in previous studies [60, 61].

According to our results, extracellular K^+ (oxygen demand) during CSD propagation, increased to identical levels under normoxia and the three pathological conditions tested, showing that oxygen demand during the tested conditions, was found identical. In contradiction, oxygen supply (CBF) and oxygen balance (mitochondrial NADH redox state) differ in amplitude values and in the simultaneous nature responses during CSD propagation, between normoxia and the pathological conditions tested (Figs. 2-7). The basic tight coupling between NADH and CBF found by the inverse oscillations between these parameters, under the pathological conditions tested, may dictate the simultaneous changes found in NADH and CBF, during CSD propagation, under hypoxia, ischemia and after L-NAME induction.

CONCLUSIONS

1. CSD induced, under conditions where oxygen delivery is reduced or disturbed, increases oxygen demand to the same level as during normoxic conditions, whereas oxygen supply (CBF) can further decrease and oxygen balance (NADH) is disturbed;
2. CSD wave duration increased under conditions where oxygen supply was reduced, indicating a reduction in energy formation;
3. The tight coupling between CBF and NADH under limited O_2 supply conditions seems to be stronger than under normoxic conditions as disclosed after increasing O_2 demand by CSD;
4. Our results demonstrate the significance of simultaneously monitoring extracellular K^+ (oxygen demand), CBF (oxygen supply) and mitochondrial NADH (oxygen balance) for reliable discernment between normal brain and conditions where oxygen balance is impaired.

CONFLICT OF INTEREST

None declared.

ACKNOWLEDGEMENT

None declared.

REFERENCES

- [1] Leao AAP. Spreading depression of activity in cerebral cortex. *J Neurophysiol* 1944; 7: 359-90.
- [2] Hansen AJ, Zeuthen T. Extracellular ion concentration during spreading depression and ischemia in the rat brain cortex. *Acta Physiol Scand* 1981; 113: 437-45.
- [3] Mayevsky A, Weiss HR. Cerebral blood flow and oxygen consumption in cortical spreading depression. *J Cereb Blood Flow Metabol* 1991; 11: 829-36.
- [4] Lauritzen M. Cerebral blood flow in migraine and cortical spreading depression. *Acta Neurolog Scand* 1987; 76: 1-40.
- [5] Somjen GG. Mechanisms of spreading depression and hypoxic spreading depression-like depolarization. *Physiol Rev* 2001; 81: 1065-96.
- [6] Somjen GG, Rosenthal M, Cordingley G, LaManna J, Lothman E. Potassium, neuroglia, and oxidative metabolism in central gray matter. *Fed Proc* 1976; 35: 1266-71.

- [7] Sonn J, Mayevsky A. Effects of brain oxygenation on metabolic, hemodynamic, ionic and electrical responses to spreading depression in the rat. *Brain Res* 2000; 882: 212-6.
- [8] Sonn J, Mayevsky A. Effects of anesthesia on the responses to cortical spreading depression in the rat brain *in vivo*. *Neurol Res* 2006; 28: 206-19.
- [9] Mayevsky A, Rogatsky G. Mitochondrial function *in vivo* evaluated by NADH fluorescence: From animal models to human studies. *Am J Physiol Cell Physiol* 2007; .292: C615-C40.
- [10] Nedergaard M, Hansen AJ. Spreading depression is not associated with neuronal injury in the normal brain. *Brain Res* 1988; 449: 395-8.
- [11] Gorji A. Spreading depression: a review of the clinical relevance. *Brain Res Brain Res Rev* 2001; 38: 33-60.
- [12] Iijima T, Mies G, Hossmann KA. Repeated negative DC deflections in rat cortex following middle cerebral artery occlusion are abolished by MK-801: effect on volume of ischemic injury. *J Cereb Blood Flow Metabol* 1992; 12: 727-33.
- [13] Hansen AJ, Nedergaard M. Brain ion homeostasis in cerebral ischemia. *Neurochem Pathol* 1988; 9: 195-209.
- [14] Mayevsky A, Doron A, Manor T, Meilin S, Zarchin N, Ouaknine GE. Cortical spreading depression recorded from the human brain using a multiparametric monitoring system. *Brain Res* 1996; 740: 268-74.
- [15] Rogatsky GG, Sonn J, Kamenir Y, Zarchin N, Mayevsky A. Relationship between intracranial pressure and cortical spreading depression following fluid percussion brain injury in rats. *J Neurotrauma* 2003; 20: 1315-25.
- [16] Gorji A, Speckmann EJ. Spreading depression enhances the spontaneous epileptiform activity in human neocortical tissues. *Eur J Neurosci* 2004; 19: 3371-4.
- [17] Cooper R, Crow HJ, Greywalter W, Winter AL. Regional control of cerebral vascular reactivity and oxygen supply in man. *Brain Res* 1966; 3: 174-9.
- [18] Takano T, Tian GF, Peng W, *et al*. Cortical spreading depression causes and coincides with tissue hypoxia. *Nat Neurosci* 2007; 10: 754-62.
- [19] Bengtsson F, Siesjo BK, Schurr A, Rigor BM. Cell damage in cerebral ischemia: Physiological biochemical and structural aspects. *Cerebral Ischemia and Resuscitation*. Boca Raton: CRC Press; 1990; pp. 215-33.
- [20] Siesjo BK, Bengtson F. Calcium fluxes, calcium antagonists, and calcium-related pathology in brain ischemia, hypoglycemia, and spreading depression: a unifying hypothesis. *J Cereb Blood Flow Metabol* 1989; 9: 127-40.
- [21] Dawson TM, Dawson VL, Snyder SH. Molecular mechanisms of nitric oxide actions in the brain. *Ann NY Acad Sci USA* 1994; 738: 76-85.
- [22] Iadecola C, Pelligrino DA, Moskowitz MA, Lassen NA. Nitric oxide synthase inhibition and cerebrovascular regulation. *J Cereb Blood Flow Metabol* 1994; 14: 175-92.
- [23] Faraci FM, Heistad DD. Regulation of the cerebral circulation: role of endothelium and potassium channels. *Physiol Rev* 1998; 78: 53-97.
- [24] Iadecola C, Zhang F, Casey R, Nagayama M, Ross ME. Delayed reduction of ischemic brain injury and neurological deficits in mice lacking the inducible nitric oxide synthase gene. *J Neurosci* 1997; 17: 9157-64.
- [25] Uetsuka S, Fujisawa H, Yasuda H, Shima H, Suzuki M. Severe cerebral blood flow reduction inhibits nitric oxide synthesis. *J Neurotrauma* 2002; 19: 1105-16.
- [26] Fujisawa H, Koizumi H, Ito H, Yamashita K, Maekawa T. Effects of mild hypothermia on the cortical release of excitatory amino acids and nitric oxide synthesis following hypoxia. *J Neurotrauma* 1999; 16: 1083-93.
- [27] Goadsby PJ, Kasube H, Hoskin KL. Nitric oxide synthesis couples cerebral blood flow and metabolism. *Brain Res* 1992; 595: 167-70.
- [28] Colonna DM, Meng W, Deal DD, Busija DW. Nitric oxide promotes arteriolar dilation during cortical spreading depression in rabbits. *Stroke* 1994; 25: 2463-70.
- [29] Sonn J, Mayevsky A. The effect of ethanol on metabolic, hemodynamic and electrical responses to cortical spreading depression. *Brain Res* 2001; 908: 174-86.
- [30] Mayevsky A. Brain NADH redox state monitored *in vivo* by fiber optic surface fluorometry. *Brain Res Rev* 1984; 7: 49-68.
- [31] Mayevsky A, Flamm ES, Pennie W, Chance B. A fiber optic based multiprobe system for intraoperative monitoring of brain functions. *Proc SPIE* 1991; 1431: 303-13.
- [32] Sokoloff L, Kenney C, Adachi K, *et al*. Effect of inhibition of nitric oxide synthase on resting local cerebral blood flow and on changes induced by hypercapnia or local functional activity. *Pharmacology of Cerebral Ischemia*. Stuttgart: Wissenschaftliche Verlagsgesellschaft mbH 1992; pp. 371-81.
- [33] Duckrow RB. A brief hypoperfusion precedes spreading depression if nitric oxide synthesis is inhibited. *Brain Res* 1993; 618: 190-5.
- [34] Faraci FM, Brain JE Jr. Nitric oxide and the cerebral circulation. *Stroke* 1994; 25: 692-703.
- [35] Sandor P, Komjati K, Nyary I. Major role of nitric oxide in the mediation of regional CO₂ responsiveness of the cerebral and spinal cord vessel of the cat. *J Cereb Blood Flow Metabol* 1994; 14: 49-58.
- [36] Macrae IM, Dawson DA, Norrie JD, McCulloch J. Inhibition of nitric oxide synthase: effects on cerebral blood flow and glucose utilisation in the rat. *J Cereb Blood Flow Metabol* 1993; 13: 985-92.
- [37] Brown GC. Nitric oxide regulates mitochondrial respiration and cell functions by inhibiting cytochrome oxidase. *FEBS Lett* 1995; 369: 136-9.
- [38] Brown GC. Regulation of mitochondrial respiration by nitric oxide inhibition of cytochrome c oxidase. *Biochim Biophys Acta* 2001; 1504: 46-57.
- [39] Kurzelewski M, Duda M, Stanley WC, Boemke W, Beresewicz A. Nitric oxide synthase inhibition and elevated endothelin increase oxygen consumption but do not affect glucose and palmitate oxidation in the isolated rat heart. *J Physiol Pharmacol* 2004; 55: 27-38.
- [40] Dirnagl U, Lindauer U, Villringer A. Role of nitric oxide in the coupling of cerebral blood flow to neuronal activation in rats. *Neurosci Lett* 1993; 149: 43-6.
- [41] Lauritzen M. Pathophysiology of the migraine aura. The spreading depression theory. *Brain* 1994; 117: 199-210.
- [42] Bredt DS, Snyder SH. Isolation of nitric oxide synthetase, a calmodulin-requiring enzyme. *Proc Natl Acad Sci USA* 1990; 87: 682-5.
- [43] Fabricius M, Akgoren N, Lauritzen M. Arginine-nitric oxide pathway and cerebrovascular regulation in cortical spreading depression. *Am J Physiol* 1995; 269: H23-H9.
- [44] Zhang F, White JG, Iadecola C. Nitric oxide donors increase blood flow and reduce brain damage in focal ischemia: evidence that nitric oxide is beneficial in the early stage of cerebral ischemia. *J Cereb Blood Flow Metabol* 1994; 14: 217-26.
- [45] Akgoren N, Fabricius M, Lauritzen M. Importance of nitric oxide for local increases of blood flow in rat cerebellar cortex during electrical stimulation. *Proc Natl Acad Sci USA* 1994; 91: 5903-7.
- [46] Dirnagl U, Niwa K, Lindauer U, Villringer A. Coupling of cerebral blood flow to neuronal activation: role of adenosine and nitric oxide. *Am J Physiol* 1994; 267: H296-H301.
- [47] Meng W, Colonna DM, Tobin JR, Busija DW. Nitric oxide and prostaglandins interact to mediate arteriolar dilation during cortical spreading depression. *Am J Physiol* 1995; 269: H176-H81.
- [48] Sukhotinsky I, Dilekoz E, Moskowitz MA, Ayata C. Hypoxia and hypotension transform the blood flow response to cortical spreading depression from hyperemia into hypoperfusion in the rat. *J Cereb Blood Flow Metabol* 2008; 28: 1369-76.
- [49] Pearce WJ. Mechanism of hypoxic cerebral vasodilation. *Pharmacol Ther* 1995; 65: 75-91.
- [50] Kuschinsky W, Wahl M. Local chemical and neurogenic regulation of cerebral vascular resistance. *Physiol Rev* 1978; 58: 656-89.
- [51] Ayata C, Shin HK, Salomone S, *et al*. Pronounced hypoperfusion during spreading depression in mouse cortex. *J Cereb Blood Flow Metabol* 2004; 24: 1172-82.
- [52] Brown GC, Borutaite V. Nitric oxide inhibition of mitochondrial respiration and its role in cell death. *Free Radic Biol Med* 2002; 33: 1440-50.
- [53] Schechter M, Sonn J, Mayevsky A. The brain oxygen balance under various experimental pathophysiological conditions. oxygen transport to tissue. *Adv Exp Med Biol* 2009; 645: 293-9.
- [54] Mayevsky A. Brain energy metabolism of the conscious rat exposed to various physiological and pathological situations. *Brain Res* 1976; 113: 327-38.
- [55] Mayevsky A. Biochemical and physiological activities of the brain as *in vivo* markers of brain pathology. In: Bernstein EF, Callow

- AD, Nicolaides AN, Shifrin EG, Eds. Cerebral, Revascularization. UK: Med Orion Pub 1993; pp. 51-69.
- [56] Hansen AJ, Schurr A, Rigor BM. Ion homeostasis in cerebral ischemia. Cerebral Ischemia and Resuscitation. Boston: CRC Press 1990; pp. 77-87.
- [57] Mayevsky A, Duckrow RB, Yoles E, Zarchin N, Kanshansky D. Brain mitochondrial redox state, tissue hemodynamic and extracellular ion responses to four-vessel occlusion and spreading depression in the rat. *Neurolog Res* 1990; 12: 243-8.
- [58] Back T, Kohno K, Hossmann KA. Cortical negative DC deflections following middle cerebral artery occlusion and KCl-induced spreading depression: effect on blood flow, tissue oxygenation and electroencephalogram. *J Cereb Blood Flow Metabol* 1994; 14: 12-9.
- [59] Mayevsky A, Zarchin N, Sonn J, Lehmenkuhler A, Grotemeyer KH, Tegtmeier F. Brain redox state and O₂ balance in experimental spreading depression and ischemia. *Migraine - Basic Mechanisms and Treatment*. Munchen-Wier, Baltimore: Urban & Schwarzenberg; 1993; pp. 379-93.
- [60] Mayevsky A, Ziv I. Oscillations of cortical oxidative metabolism and microcirculation in the ischemic brain. *Neurolog Res* 1991; 13: 39-47.
- [61] Meilin S, Mendelman A, Sonn J, Manor T, Zarchin N, Mayevsky A. Metabolic and hemodynamic oscillations monitored optically in the brain exposed to various pathological states. Oxygen transport to tissue. *Adv Exp Med Biol* 1999; 471: 141-6.

Received: December 02, 2011

Revised: January 14, 2012

Accepted: January 18, 2012

© Sonn and Mayevsky; Licensee *Bentham Open*.

This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.